

RECOMMENDED MINIMUM EXCLUSION PERIODS

CONDITION	EXCLUSION	EXCLUSION OF CONTACTS
Campylobacter infection	Exclude until there has not been a loose bowel motion for 24 hours ^b	Not excluded
Candidiasis (thrush)	Not excluded	Not excluded
Cytomegalovirus (CMV) infection	Not excluded	Not excluded
Conjunctivitis	Exclude until discharge from the eyes has stopped, unless a doctor has diagnosed non-infectious conjunctivitis	Not excluded
Cryptosporidium	Exclude until there has not been a loose bowel motion for 24 hours ^b	Not excluded
Diarrhoea (No organism identified)	Exclude until there has not been a loose bowel motion for 24 hours ^b	Not excluded
Fungal infections of the skin or nails (e.g. ringworm, tinea)	Exclude until the day after starting appropriate antifungal treatment	Not excluded
Giardiasis	Exclude until there has not been a loose bowel motion for 24 hours ^b	Not excluded
Glandular fever (mononucleosis, Epstein Barr virus [EBV] infection)	Not excluded	Not excluded
Hand, foot and mouth disease	Exclude until all blisters have dried	Not excluded
Haemophilus influenzae type b (Hib)	Exclude until the person has received appropriate antibiotic treatment for at least 4 days	Not excluded. Contact a public health unit for specialist advice
Head lice (pediculosis)	Not excluded if effective treatment begins before the next day at the education and care service. The child does not need to be sent home immediately if head lice are detected	Not excluded
Hepatitis A	Exclude until a medical certificate of recovery is received and until at least 7 days after the onset of jaundice	Not excluded. Contact a public health unit for specialist advice about vaccinating or treating children in the same room or group
Hepatitis B	Not excluded	Not excluded
Hepatitis C	Not excluded	Not excluded
Herpes simplex (cold sores, fever blisters)	Not excluded if the person can maintain hygiene practices to minimise the risk of transmission. If the person cannot comply with these practices (e.g. because they are too young), they should be excluded until the sores are dry. Sores should be covered with a dressing, where possible	Not excluded
Human immunodeficiency virus (HIV)	Not excluded. If the person is severely immune compromised, they will be vulnerable to other people's illnesses	Not excluded
Hydatid disease	Not excluded	Not excluded
Impetigo	Exclude until appropriate antibiotic treatment has started. Any sores on exposed skin should be covered with a watertight dressing	Not excluded
Influenza and influenza-like illnesses	Exclude until person is well.	Not excluded
Listeriosis	Not excluded	Not excluded
Measles	Exclude for at least four days after the onset of rash.	Immunised and immune contacts are not excluded For non-immunised contacts, contact a public health unit for specialist advice. All

		immunocompromised children should be excluded until 14 days after the appearance of the rash in the last case
Meningitis (viral)	Exclude until person is well.	Not excluded
Meningococcal infection	Exclude until appropriate antibiotic treatment has been completed	Not excluded. Contact a public health unit for specialist advice about antibiotics and/or vaccination for people who were in the same room as the case
Molluscum contagiosum	Not excluded	Not excluded
Mumps	Exclude for 9 days or until swelling goes down (whichever is sooner)	Not excluded
Norovirus	Exclude until there has not been a loose bowel motion or vomiting for 48 hours	Not excluded
Pertussis (whooping cough)	Exclude until 5 days after starting appropriate antibiotic treatment, or for 21 days from the onset of coughing	Contact a public health unit for specialist advice about excluding non-vaccinated contacts, or antibiotics
Pneumococcal disease	Exclude until person is well	Not excluded
Roseola	Not excluded	Not excluded
Ross River virus	Not excluded	Not excluded
Rotavirus infection	Exclude until there has not been a loose bowel motion or vomiting for 24 hours ^b	Not excluded
Rubella (German measles)	Exclude until fully recovered or for at least 4 days after the onset of the rash	Not excluded
Salmonellosis	Exclude until there has not been a loose bowel motion for 24 hours ^b	Not excluded
Scabies	Exclude until the day after starting appropriate treatment	Not excluded
Shigellosis	Exclude until there has not been a loose bowel motion for 24 hours ^b	Not excluded
Streptococcal sore throat (including scarlet fever)	Exclude until the person has received antibiotic treatment for at least 24 hours and feels well	Not excluded
Toxoplasmosis	Not excluded	Not excluded
Tuberculosis (TB)	Exclude until medical certificate is produced from the appropriate health authority	Not excluded. Contact a public health unit for specialist advice about screening, antibiotics or specialist TB clinics
Varicella (chickenpox)	Exclude until all blisters have dried—this is usually at least 5 days after the rash first appeared in non-immunised children, and less in immunised children	Any child with an immune deficiency (for example, leukaemia) or receiving chemotherapy should be excluded for their own protection. Otherwise, not excluded
Viral gastroenteritis (viral diarrhoea)	Exclude until there has not been a loose bowel motion for 24 hours ^b	Not excluded

The definition of 'contacts' will vary according to the disease—refer to the specific fact sheet for more information.^b If the cause is unknown, possible exclusion for 48 hours until cause is identified. However, educators and other staff who have a food handling role should always be excluded until there has not been a loose bowel motion for 48 hours. Adapted from SA Health Communicable Disease Control Branch: <http://www.dh.sa.gov.au/pehs/branches/branch-communicable.htm>. Note that exclusion advice is consistent

with Series of National Guidelines (SoNGs) where available. Staying Healthy in Childhood education and care. Preventing infectious diseases | 5th Edition | ref # CH000a